



Assessing vulnerability and adaptation of river systems

Massimo Menenti
Delft University of Technology




Climate Change Adaptation

- hydrological observations, analyses and predictions
- mitigate the impacts of climate change
- remove the underlying causes of vulnerability to climate change
- Themes:
 - Short- and long-term hydrological variability and climate change
 - Resilience versus vulnerability of hydrologic systems
 - Resilience of hydrological systems to climate variability
 - Identifying and removing the causes of the vulnerability of hydrological systems to climate change
 - Reconstruction of adaptation mechanisms to hydrological variability in the recent past
- GEWEX / HAP Working Group?
- Need to integrate very diverse data and models river-wise



Water Balance Model domain





CEOP-AEGIS



EU Project

Collaborative
Project / Small or medium-scale focused research project –
Specific International Co-operation Action



ENV.2007.4.1.4.2
Improving observing systems for
water resource management

Period: May 2008 – May 2012

Coordinated Asia-European long-term Observing system of Qinghai-Tibet Plateau hydro-meteorological processes and the Asian-monsoon system with Ground satellite Image data and numerical Simulations

An international cooperation project between Europe and Asia to improve knowledge on hydrology and meteorology of the Tibetan Plateau and its role in climate, monsoon and extreme meteorological events

- To construct out of existing ground measurements and current / future satellites an observing system to determine and monitor the water yield of the Plateau, i.e. how much water is finally going into the seven major rivers of SE Asia
- To monitor the evolution of snow, vegetation cover, surface wetness and surface fluxes and analyze the linkage with convective activity, (extreme) precipitation events and the Asian Monsoon.

19 beneficiaries from 8 european and asian countries




www.ceop-aegis.org





CEOP AEGIS concept





Climate elasticity of streamflow

(Zheng et al., 2009)

$$Q = f(P, E_0, V)$$

Q = streamflow

P = precipitation

E_0 = potential evapotranspiration

V = land use / land cover

$$\Delta Q = f'_P \Delta P + f'_{E_0} \Delta E_0 + f'_V \Delta V$$

$$\varepsilon = \frac{dQ / Q}{dX / X}$$

climate elasticity of streamflow to X


case study on the Headwaters of the Yellow River Basin (HYRB)

land use change: about 74.6% of the streamflow reduction in the 1990s

climate change 25.4% of the reduction




Observed trends Qinghai – Tibet Plateau 1960 - 2000






Potential ET









Precipitation






Lake level changes from ICESAT/GLAS 2003 - 2009

(Phan Hien et al., 2010)





Regional variability of trends






Number of frozen days in China (Oct. 1, 2002 ~ Sep. 31, 2003)


**Sensor:
SSM/I**

**Method:
Decision
Tree**





ET deficit year 2009





PBL response to land surface forcing



GRAPES = Global / Regional Assimilation and PrEdiction System

unified NWP model with 3/4DVAR data assimilation system under development by Shen et al. at CMA / CAMS, China.


GRAPES_Meso with 15 km x 15 km horizontal resolution

Dynamics	Flux-form equations of water substances
	Piece-wise rational method + volume-remapping SL for scalar advection
	Reference atmosphere based on the initial field
	Effective topography
Physics	NOAH LSM + Simple initialization
	Xu and Randall Diagnostic cloud
	Betts-Miller-Janjic Cumulus
	CAMS mixed-phase microphysics
	Effect of slope on surface radiation




Describing the ABL of the Plateau

30km simulation, 30' step, 48h runs – Ta, q, u, v, p, Global, IRa






ABL height daily cycle





Describing the ABL of the Plateau





Generalize the Energy Balance mapping on the Plateau

- SEBI based (Menenti and Choudhury, 1993)
- Physical calculation of the extreme boundaries following the SEBS approach (Su, 2000)
- 2D ABL forcing & calculation grid size for stability functions depending on the ABL development (MSSEBS, Colin 2006)
- Expected results:
 - **1km resolution Surface Energy Balance** components
 - **Time series** on a week to 10 days basis
 - Automated processing chain, including interface with data providers & results repositories

Meso-scale Atmospheric Forcing grid [15-100 km]


T_a, q, u, v, p

Sw, Lw incoming radiance

ABL Calculation grid [10 x \overline{ABL} height]


Full resolution calculation grid [TIR Resolution]

$LST, albedo, fc, LAI, emissivity, DEM$






Spatial and temporal variability PBL height





PBL Height (m) – GRAPES Meso

14 January 08




14 July 08






Plateau Water Balance: Data Requirements








First results: routed discharge



Immerzeel et al., 2010. Science



Large fluctuations of regional and local water cycles





Climate Change Will Affect the Asian Water Towers


Immerzeel et al., 2010



The map shows the importance of melt water to downstream areas (pie charts) and the number of people threatened in their food security due to reduction in upstream flow due to climate change




Daily streamflow





CEOP-AEGIS Database

- Principles
 - all project products in one single entry point
 - web-based user-friendly interface
 - registered as a contribution to GEOSS
 - open access to the scientific community





List of datasets

- D1.4 Final 3 years dataset on aerodynamic and thermodynamic variables over the different land surfaces of the Tibetan Plateau
- D2.5 Data set vegetation cover, surface albedo, LAI, aerosol optical depth, precipitable water vapour, soil/foilage component and land surface emissivity/temperatures for a 3 years period.
- D3.2 Generalize SEB calculation at a high spatial resolution and on a regional extent
- D3.4 Validated complete time-series maps of fluxes over 3 years, with a target frequency of one week
- D4.1 Collection of consistent continuous in-situ soil moisture measurements at regional scale
- D4.3 Estimation of soil moisture from Geostationary Satellite (GS) data
- D4.4 A data product of the plateau using different sensors simultaneously
- D4.5 Validation results and documentation of uncertainties
- D5.3 Provide 3-years 3-D gridded radar data for case studies
- D5.4 Provide high space and temporal resolution precipitation data retrieved from radar, satellite (both VIS-IR and MW) and ground rain gauge data for case studies
- D6.4 Validated data set (include snow cover area, SWE, glacier area and storage from satellite data, and assimilation data sets) from 2008 to 2010
- D8.2 Time series analysis of water balance data
- D9.1 Vegetation dynamic maps over a long-term period
- D9.2 Maps of drought vulnerability zones in the study area
- D9.5 Anomalies maps of rainfall, vegetation response
- D10.3 Flood inundation, flood hazard and flood risk zone maps based on the hydraulic modelling and analysis of satellite data
- D10.4 Real time forecasting of flood inundation, depth, areal extent and return period of flooding
- D10.5 Case studies on flood events in the period 1999-2010



List of datasets partially or totally available


- D2.5 Data set vegetation cover, surface albedo, LAI, aerosol optical depth, precipitable water vapour, soil/foilage component and land surface emissivity/temperatures for a 3 years period.
- D3.2 Generalize SEB calculation at a high spatial resolution and on a regional extent
- D4.1 Collection of consistent continuous in-situ soil moisture measurements at regional scale
- D4.3 Estimation of soil moisture from Geostationary Satellite (GS) data
- D4.4 A data product of the plateau using different sensors simultaneously
- D5.3 Provide 3-years 3-D gridded radar data for case studies
- D5.4 Provide high space and temporal resolution precipitation data retrieved from radar, satellite (both VIS-IR and MW) and ground rain gauge data for case studies
- D6.4 Validated data set (include snow cover area, SWE, glacier area and storage from satellite data, and assimilation data sets) from 2008 to 2010
- D8.2 Time series analysis of water balance data
- D9.1 Vegetation dynamic maps over a long-term period
- D9.2 Maps of drought vulnerability zones in the study area
- D9.5 Anomalies maps of rainfall, vegetation response

CEOP-AEGIS Data Repository
Hydrology and Climatology of the Tibetan Plateau


Center map at location:

Drag a region to plot with mouse

Help Hide map



Hydrology
PCRGLOB-WB Model 2008



Plot Selected

Link to this page

Datasets

Search

Category	Dataset
Atmosphere	Global Reanalysis
Atmosphere	Regional Low-resolution Reanalysis
Hydrology	PCRGLOB-WB Model 2008
Hydrology	PCRGLOB-WB Model 2009

Variables

Search

specific runoff

Plot type

- Shaded 2D plot
- Filled 2D contour plot
- 1D Line plot
- 1D Bar plot
- Animate in Google Earth
- Download data

Ranges

Lon

Lat

Time

Reset

Plot view

- Lon/Lat(xy)
- Lon/Time(xt)
- Lat/Time(yt)

Plot options

Auto colorbar:

Colorbar range:

Plot selected Link to this page


Done

DChart - CEOP-AEGIS Data Repository - Hydrology and Climatology of the Tibetan Plateau - Mozilla Firefox

File Edit View History Bookmarks Tools Help

http://localhost:8080/dchart/index.html?dsetid=bec8437c48ade3ac9410081ea74146a&yv test

LOCATION MAP



DATASETS


Category	Dataset
Atmosphere	Global Reanalysis
Atmosphere	Regional Low-resolution Reanalysis
Hydrology	PCRGLOB-WB Model 2008
Hydrology	PCRGLOB-WB Model 2009

VARIABLES

specific runoff

Visualisation

PCRGLOB-WB Model 2008
Time: 01 Aug 2008 00:00:00 GMT



specific runoff(meters per day)

Plotting tools

- Filled 2D contour plot
- 1D Line plot
- 1D Bar plot
- Animate in Google Earth
- Download data

EXPORT

Plot view

- Lon/Lat(xy)
- Lon/Time(xt)
- Lat/Time(yt)

Plot options

Auto colorbar:

Colorbar range:

Space & time

Lon:

Lat:

Time:

Reset

Plot selected Link to this page

Done



Data analysis and integration

- Merge radiometric data optical systems
- Cross calibration “family” imaging radiometers
- Develop generic algorithms
- Analysis of time series of satellite observations to get statistics on anomalies (when, where, how large) on droughts and floods
- 30+ years AVHRR , microwave radiometers
- Access to data bases of different initiatives: Third Pole Environment (China, Nepal...), PAPRIKA (Pakistan, Italy, France)



Acknowledgements



- This study is jointly supported by the **ESA Dragon II** program under proposal no.5322: “Key Eco-Hydrological Parameters Retrieval and Land Data Assimilation System Development in a Typical Inland River Basin of China’s Arid Region” and by the **European Commission** (Call FP7-ENV-2007-1 Grant nr. 212921) as part of the **CEOP-AEGIS** project (<http://www.ceop-aegis.org>) coordinated by the University of Strasbourg.
- **WATER** is jointly supported by the Chinese Academy of Science Action Plan for West Development Program (grant KZCX2-XB2-09) and Chinese State Key Basic Research Project (grant 2007CB714400).